

PROCEEDINGS

OF THE

SCIENTIFIC MEETINGS

OF THE

ZOOLOGICAL SOCIETY

OF LONDON

FOR THE YEAR

1869.

PRINTED FOR THE SOCIETY,
AND SOLD AT THEIR HOUSE IN HANOVER SQUARE.

LONDON:
MESSRS. LONGMANS, GREEN, READER, AND DYER,
PATERNOSTER ROW.

LIST
OF THE
CONTRIBUTORS,

With References to the several Articles contributed by each.

	Page
ADAMS, HENRY, F.L.S.	
Descriptions of a New Genus and Fourteen New Species of Marine Shells. (Plate XIX.)	272
ALLPORT, MORTON, F.Z.S.	
Letter from, on the Introduction of Salmon into Tasmania .	473
ANDERSON, Dr. JOHN, C.M.Z.S.	
Letter from, relating to his travels in Yunan and Upper Burmah	111
Letter from, concerning his recent Expedition to Yunan, and Animals destined for the Society's Menagerie	277
Letter from, relating to living Specimens of <i>Ailurus fulgens</i>	278
Letter from, relating to Indian Animals (<i>Porcula salvania</i> and <i>Budorcas taxicolor</i>)	470
ANGAS, GEORGE FRENCH, F.L.S., C.M.Z.S., &c.	
Descriptions of Twelve New Species of Land and Marine Shells from Australia and the Solomon Islands. (Plate II.)	45
Descriptions of Eight New Species of <i>Helicidæ</i> from the Western Pacific Islands. (Plate XLVIII.)	624

LIST OF PLATES.

Plate		Page
I.	<i>Gonostomyx loa-loa</i>	38
II.	New Shells from Australia and the Solomon Islands ..	45
III.	} Siliceo-fibrous Sponges	66
IV.		
V.		
VI.		
VII.	<i>Otaria jubata</i>	100
VIII.	<i>Embletonia grayii</i>	109
IX.	<i>Gracula krefftii</i>	} 118
X.	<i>Rallus intactus</i>	
XI.	<i>Domicella cardinalis</i>	126
XII.	<i>Neorhynchus nasesus</i> *	146
XIII.	<i>Poospiza cæsar</i>	151
XIV.	<i>Chatops grayii</i>	163
XV.	{ Fig. 1. <i>Trionyx formosus</i>	} 165
	{ Fig. 2. <i>Baikiea elegans</i>	
XVI.	<i>Chatodon dichrous</i>	238
XVII.	Nests of <i>Ploceus baya</i>	243
XVIII.	<i>Brachygalba goeringi</i>	250
XIX.	New Marine Shells.....	272
XX.	<i>Phacochærus æliani</i> , ♀	276
XXI.	} Siliceo-fibrous Sponges	323
XXII.		
XXIII.		
XXIV.		
XXV.		
XXVI.	New Australian Shells	358
XXVII.	Eggs of Australian Cuckoos and their Foster-parents ..	359
XXVIII.	<i>Calliste florida</i>	416
XXIX.	<i>Buccanodon anchietæ</i>	436
XXX.	{ Figs. 1, 2. <i>Euphonia chrysopasta</i> , ♂ et ♀	} 437
	{ Fig. 3. <i>Neopipo rubicunda</i>	
XXXI.	<i>Buthraupis arcæi</i>	} 439
XXXII.	<i>Tachyphonus chrysomelas</i>	

* Err. *maseus* on Plate.

4. Descriptions of a New Genus and Fourteen New Species of Marine Shells. By HENRY ADAMS, F.L.S.

(Plate XIX.)

CONUS (CORONAXIS) CERNICUS, Barcl. MS. (Plate XIX. fig. 1.)

C. testa conica, solida, polita, spiraliter puncto-striata, striis antice validioribus et propinquieribus; spira elevata, convexo-conica, sutura valde impressa; anfr. 9, angulatis, coronatis; cinereo-alba, rubido-fulvo fasciata et nebulosa, maculis albis aspersis, juxta suturas rubido-fulvo notata; apertura angusta, lineari.

Long. 25, lat. 12 mill.

Hab. Barkly Island, Mauritius (*Coll. Barclay*).

This and several of the following species were procured at Mauritius after a late severe storm, by which large masses of coral were displaced and formed into a small island on the coast, which has been named Barkly Island, after the Governor Sir Henry Barkly.

CLATHURELLA ROBILLARDI, Barcl. MS. (Plate XIX. fig. 2.)

C. testa turrita, tenui, alba, costis validis rotundatis ad 10 longitudinaliter ornata, costulis transversis et striis longitudinalibus cancellata; anfr. 9, convexis, postice excavatis, ultimo $\frac{1}{3}$ longitudinis testæ; apertura anguste lunata; columella costulis obliquis insculpta; labro incrassato, intus subsulcato; sinu angusto, haud profundo; rostro longo, recurvo.

Long. 25, lat. $8\frac{1}{2}$ mill.

Hab. Barkly Island, Mauritius (*Coll. Barclay*).

DRILLIA BARKLIENSIS, H. Ad. (Plate XIX. fig. 3.)

D. testa elongato-turrita, roseo-fulvida, longitudinaliter striata, ad basim plicata; spira anfr. 8, prope suturam excavatis, costulis granulosis (granulis albis) cinctis, interstitiis transversim striatis, ad medium nodulis albis ornatis; apertura oblonga, $\frac{1}{3}$ longitudinis testæ aequante; columella rectiuscula; sinu mediocri; labro extus varicoso, macula fulva notato.

Long. 18, lat. $6\frac{1}{2}$ mill.

Hab. Barkly Island, Mauritius (*Coll. H. Ad.*).

I am indebted to Mr. Caldwell, of Mauritius, for an example of this species.

CORALLIOPHILA CORONATA, Barcl. MS. (Plate XIX. fig. 4.)

C. testa imperforata, ovato-fusiforimi, plicis longitudinalibus distantibus, et costis transversis undulatis confertis sculpta, in medio anfractus ultimi costis duabus validioribus, posteriore spinis curvatis armata; purpureo-fulva, costis transversis pallidioribus; anfr. 7, convexis, ultimo dimidiam longitudinis testæ superante; apertura acuminato-ovata, intus violacea; columella levi, sim-

HAMINEA SUBPELLUCIDA, H. Ad. (Plate XIX. fig. 13.)

H. *testa tenuissima, albida, subpellucida, ovata, in medio plerumque gibbosa, longitudinaliter striis undulatis rugosis, et spiraliter lineis elevatis irregularibus sculpta; vertice excavato, subperforato; apertura antice dilatata; margine columellari simplici, arcuato; labro recto, postice rotundato.*

Long. 17, lat. 11 mill.

Hab. Lisbon (Coll. M'Andrew).

GOULDIA MODESTA, H. Ad. (Plate XIX. fig. 14.)

G. *testa subtriangulari, solidula, costis concentricis insculpta, intervalis costis aequalibus, pallide fusca, maculis rubris paucis picta; extremitate antica versus umbones concava, postica arcuata; umbonibus acutis approximatis; margine ventrali convexo, intus crenulato.*

Long. $6\frac{1}{2}$, alt. 6 mill.

Hab. Gulf of Tunis (Coll. M'Andrew).

I take this opportunity of correcting a few names given by me to shells in former papers, which I have since found to be preoccupied:—

PLANORBIS (ADULA) SEPTEMVOLVIS to be altered to PLANORBIS (ANÆUS) SEPTEMVOLVIS (Proc. Zool. Soc. 1861, p. 145).

HELIX (GEOTROCHUS) BLANFORDI to be altered to HELIX (GEOTROCHUS) BLANFORDIANA (Proc. Zool. Soc. 1865, p. 415).

PUPINA PFEIFFERI to be altered to PUPINA PFEIFFERIANA (Proc. Zool. Soc. 1865, p. 416).

STYLODONTA (EREPTA) RUFOCINCTA to be altered to STYLODONTA (EREPTA) RUFOZONATA (Proc. Zool. Soc. 1867, p. 303).

NANINA (?ROTULA) CONULUS to be altered to NANINA (?ROTULA) TURRITELLA (Proc. Zool. Soc. 1867, p. 307).

DESCRIPTION OF PLATE XIX.

- Fig. 1. *Conus (Coronaxis) cernicus*, p. 272.
 2. *Clathrella robillardi*, p. 272.
 3. *Drillia barkliensis*, p. 272.
 4. *Coralliophila coronata*, p. 272.
 5. 5a. *Mauritia barclayi*, p. 273.
 6. 6a. *Marginella (Glabella) mirabilis*, p. 273.
 7. 7a. *Nacella (Cellana) cernica*, p. 273.
 8. *Natica marmorata*, p. 274.
 9. *Scala delicatula*, p. 274.
 10. *Syrnola minuta*, p. 274.
 11. *Turbonilla speciosa*, p. 274.
 12. *Cancellaria pusilla*, p. 274.
 13. *Haminea subpellucida*, p. 275.
 14. *Gouldia modesta*, p. 275.

