

VOL. 1.

NO. 3.

FOSSIL SHELLS

OF

THE TERTIARY FORMATIONS

OF

NORTH AMERICA.

ILLUSTRATED BY FIGURES DRAWN ON STONE,
FROM NATURE.

BY T. A. CONRAD,

Member of the Acad. Nat. Sc. of Philada.

CONTENTS.

| | | |
|---------------------|-------------------------|------------------------|
| Voluta Sayana. | Infundibulum urticosum. | Siliquaria vitis. |
| Voluta petrosa. | Pyrula penita. | Cytherea Poulsoni. |
| Fusus trabeatus. | Fissurella tenebrosa. | Cytherea Hydana. |
| Fusus papillatus. | Conus sauridens. | Cytherea æquorea. |
| Fusus inauratus. | Marginella larvata. | Cytherea discoidalis. |
| Fusus thoracicus. | Marginella crassilabra. | Cytherea perovata. |
| Melongena armigera. | Dentalium thalloides. | Nucula magnifica. |
| Murex engonatus. | Volvaria galba. | Arca cuculoides. |
| Scalaria nassula. | Mitra bolaris. | Corbula nasuta. |
| Solarium alveatum. | Mitra doliata. | Astarte callosa. |
| Solarium antrosum. | Buccinum sagenum. | Astarte proruta. |
| Rostellaria velata. | Cancellaria gemmata. | Plicatula filamentosa. |
| Oliva Alabamensis. | Littorina antiquata. | |
| Oliva bombylis. | Melania vetusta. | |

PHILADELPHIA :

Sold by JUDAH DOBSON, No. 108, Chestnut St.

August, 1833.

W. P. Gibbons, Pr. cor. Sixth and Cherry Sts.

body whorl: striæ alternating in size: five volutions; spire elevated and pointed. Lip thickened towards the margin.

Variety, C. Destitute of carinations, and probably the young shell. Length about an inch.

Locality, Claiborne, Alab.

Cab. Acad. N. S.

FISSURELLA.

F. tenebrosa. Pl. 17. fig. 5.

Shell ovate or oval, elevated, with numerous delicate raised costæ, decussated by transverse striæ; inferior margin with impressed lines: apex pointed. Length 1 inch.

Locality, Claiborne, Alab.

Cab. Acad. N. S.

CONUS.

C. sauridens. Pl. 17. fig. 6.

Shell smooth except at base, which is obliquely striated; whorls of the spire flattened, striated; apex pointed; mouth narrow; shoulder angular. Length 1 inch.

Locality, Claiborne, Alab.

Cab. Acad. N. S.

MARGINELLA.

1. *M. larvata*. Pl. 17. fig. 7.

Shell smooth, thick; spire obsolete; with from six to eight teeth on the columella. Length $\frac{3}{4}$ of an inch.

Locality, Claiborne, Alab.

Cab. Acad. N. S.

2. *M. crassilabra*. Pl. 17. fig. 8.

Shell smooth, thick; spire elevated; varix on the outer lip very thick and extending upon the spire; lip crenated on its inner edge; nine or ten teeth on the columella. Length less than $\frac{1}{2}$ an inch.

Locality, Claiborne, Alab.

Cab. Acad. N. S.