

Description of NEW SPECIES OF SHELLS, by JOHN C. JAX, M. D.

Read February 1, 1846.

BULIMUS INFUNDIBULIFORMIS. Plate X. fig. 7, *a. b.*

Bul. testâ conicâ, tenui, griseâ, longitudinaliter striatâ, umbilico magno, infundibuliformi; spirâ conicâ, elongatâ; apice acuto; anfractibus septem subplanis; suturâ profundâ; aperturâ ovali, albidâ; labro acuto.

Shell conical, thin, gray, longitudinally striated, umbilicus very large, and funnel-shaped; spire conical and elongated; apex acute; whorls seven and somewhat flattened, sutures profound; aperture oval, rather compressed, whitish; lip acute: length, 1-2 inch, breadth, 5-16 inch.

Habitat. Interior of Bolivia.

CONUS CAILLIAUDII. Plate X. fig. 8, *a. b.*

Con. testâ oblongo-turbanatâ, subfusiformi; fuscâ, filis tenuissimis cinctâ; spirâ turrito-exsertâ; apice valdè elato; labro tenui, acuto, arcuato, juxtâ spiram emarginato; aperturâ fauce fuscâ.

Shell oblong turbinated, somewhat fusiform, brown, encircled with a number of very fine thread-like lines; spire exserted in the form of a turret, apex very elevated; lip thin, sharp, arched, emarginated next the spire, aperture brown with a white blotch: length, 1 5-8 inch, breadth, 4-8 inch.

Habitat?

The remarkable feature of this shell is the spire, which is exactly half its length, the whorls very prominent and nine in number.

I am indebted to Mons. F. Cailliaud of Nantes, for this rare Cone, and have taken the liberty to dedicate it to him.

I embrace this opportunity to make several corrections in the last Edition of the Catalogue of my Cabinet.

Plate 1, fig. 1. *Bulimus cinctus*, Nobis, is *Bulimus Favannii*, Lam. Delessert, plate 27, fig. 8.

Habitat, St. Augustine Bay, Madagascar.

“ “ *fig. 2, 3.* *Turbo rotelliformis*, nobis, is *Trochiscus Norrissii*, Sowerby Beechey Zool. plate 34, fig. 14.

“ “ *fig. 4.* *Ampullaria Brownii*, nobis, is *A. crassa*, Swains. var.

“ “ *fig. 5.* *Ampullaria Storeria*, nobis, is *A. naticoides*, D’Orbigny.

“ “ *fig. 6, 7.* *Helix Planorbis*, nobis, is *H. monilis*, Brod, Zool. P., 1832.

“ “ *fig. 8, 9.* *Physa scalaris*, nobis, see Haldeman’s Monograph, page 34, plate 4, fig. 9.

“ “ *fig. 10, 11.* *Lymnea gracilis*, nobis, see Hald. Mon. page 50, plate 13, fig. 21.

Plate 6, fig. 1. *Bulimus ustulatus*, nobis, is a variety of *Bul. chrysalidiformis*, Sowb. Zool. Proc.

Plate 7, fig. 1. *Paludina tristis*, nobis, is *P. olivacea*, Sowb.

“ “ *fig. 2, 3.* *Conus rhododendron*, Couthouy, is *Conus Adamsoni*, Gray.

“ “ *fig. 4, 5.* *Cyclostoma cumingii*, Sowb. is *C. gigantea*, Sowb.

Plate 10, fig. 1, 2. *Voluta armata?* Lam. var. is *V. Miltoni*, Gray.

ERRATA.

Through an oversight the following errata have occurred :

In pages 142 to 156, for Tab. I., read Tab. VII., for Tab. II., read Tab. VIII., and for Plate III., read Plate IX., wherever occurring.

Fig. 6.




Fig. 6. b.



Fig. 7.



Fig. 6. c.



Fig. 8.




Fig. 2.



b.



Fig. 3.



b.



Fig. 5.



Fig. 7

